arXiv:0709.3227v2 [hep-th] 15 Feb 2008

Green functions and dimensional reduction of quantum fields on product manifolds

Z. Haba
Institute of Theoretical Physics, University of Wroclaw,
50-204 Wroclaw, Plac Maxa Borna 9, Poland
e-mail:zhab@ift.uni.wroc.pl

Abstract

We discuss Euclidean Green functions on product manifolds $\mathcal{P} = \mathcal{N} \times \mathcal{M}$. We show that if $\mathcal{M}$ is compact and $\mathcal{N}$ is not compact then the Euclidean field on $\mathcal{P}$ can be approximated by its zero mode which is a Euclidean field on $\mathcal{N}$. We estimate the remainder of this approximation. We show that for large distances on $\mathcal{N}$ the remainder is small. If $\mathcal{P} = R^{D-1} \times S^{\beta}$, where S^{β} is a circle of radius β, then the result reduces to the well-known approximation of the D dimensional finite temperature quantum field theory by $D-1$ dimensional one in the high temperature limit. Analytic continuation of Euclidean fields is discussed briefly.

1 Introduction

The aim of the Kaluza-Klein program [1] is a unification of interactions by means of an extension of the number of dimensions D. Then, $D-4$ dimensions are supposed to be unobservable. In this paper we ask the question whether a higher dimensional quantum field theory can be approximated by a lower dimensional one (not necessarily from the point of view of the Kaluza-Klein program). We have shown in our earlier paper [2] (see also [3]) that such an approximation applies near the bifurcate Killing horizon when D dimensional quantum field theory can be approximated by a two-dimensional one. We are interested in the dimensional reduction from the point of view of Green functions,i.e., correlation functions of quantum fields. We examine the question whether the compact dimensions are negligible. In terms of the Green functions this means that Green functions in higher dimensions should be approximated by the ones in lower dimensions. Such a property cannot be true at arbitrarily small distances (except of some brane-type models [4][5]) because the singularity of the Green function depends on the dimension. So, the approximation can make sense only above a certain length scale. If the manifold $\mathcal{P}$ is the product manifold

$\mathcal{P}=\mathcal{N}\times\mathcal{M}$ then we expect that the Green functions on $\mathcal{P}$ can be approximated by the ones on $\mathcal{N}$ if the distances on $\mathcal{N}$ are big in comparison to the size of $\mathcal{M}$. In the conventional approach to Kaluza-Klein theories [1], the Fourier expansion of functions on the compact manifold $\mathcal{M}$ leads to large masses which make (by a formal argument) propagators of the massive Kaluza-Klein particles negligible, realizing in this way the dimensional reduction. We are interested to see how this decoupling works in the configuration space.

In order to study the problem in a rigorous way we work in the Euclidean (Riemannian instead of pseudoRiemannian) formulation of quantum field theory. Although the Euclidean approach to fields on a manifold is not as complete as in the flat space there are already some crucial results concerning the analytic continuation and construction of quantum fields [6][7]. The quantum fields are determined by the Green functions. We discuss in this paper only the two-point function which is sufficient for a construction of free fields. The analytic continuation of interacting Euclidean fields can be performed if $\mathcal{P}$ and the interaction have an additional reflection symmetry [6][7].

In sec.2 we define the Green functions. The Green functions are expanded in eigenfunctions in sec.3. We distinguish the contribution of the zero mode which determines the dimensional reduction. In sec.4 we discuss a special case of two dimensional manifold $\mathcal{N}$.We estimate the correction to the lower dimensional approximation in sec.5. In sec.6 we construct quantum fields from the Green functions. We discuss a possible extension of the results in sec.7.

2 Warped metric on a product manifold

We consider a manifold in the form of a product $\mathcal{P}=\mathcal{N}\times\mathcal{M}$ where $\mathcal{N}$ has $D-d$ dimensions and $\mathcal{M}$ is a d dimensional manifold . We assume that a metric on $\mathcal{P}$ can be expressed in the warped form [8]

$$ds^2=\sigma_{AB}dX^AdX^B=g_{ab}(x)dx^adx^b+v^2(x)h_{jk}(y)dy^jdy^k \tag{1}$$

where the coordinates on $\mathcal{P}$ are denoted by the capital $X=(x,y)$, the ones on $\mathcal{N}$ by x and the coordinates on $\mathcal{M}$ are denoted by y. If $v=1$ then the metric on the product manifold is just a product of the metrics.

Let

$$\triangle_P=\frac{1}{\sqrt{\sigma}}\partial_A\sigma^{AB}\sqrt{\sigma}\partial_B \tag{2}$$

be the Laplace-Beltrami operator on $\mathcal{P}$. We are interested in the calculation of the Green functions ($\sigma=\det(\sigma_{AB})$)

$$(-\triangle_P+m^2)\mathcal{G}^m=\frac{1}{\sqrt{\sigma}}\delta \tag{3}$$

In the metric (1) eq.(3) reads

$$\Big(-v^{d-2}\sqrt{g}(\triangle_M-m^2)-\partial_ag^{ab}v^d\sqrt{g}\partial_b\Big)\mathcal{G}^m=h_M^{-\frac{1}{2}}\delta(X-X') \tag{4}$$

A solution of eq.(3) can be expressed by the fundamental solution of the diffusion equation

$$\partial_\tau P_\tau = \frac{1}{2}\triangle_P P_\tau \tag{5}$$

with the initial condition $P_0(X,X') = \sigma^{-\frac{1}{2}}\delta(X-X')$. Then

$$\mathcal{G}^m = \frac{1}{2}\int_0^\infty d\tau \exp(-\frac{1}{2}m^2\tau)P_\tau \tag{6}$$

If $v=1$ then we may write eq.(4) in the form (here $h_M = \det(h_{jk})$ and $g = \det(g_{ab})$)

$$(-\triangle_M + m^2 - \triangle_N)\mathcal{G}^m = h_M^{-\frac{1}{2}}g^{-\frac{1}{2}}\delta(X-X') \tag{7}$$

In such a case from eq.(5) we obtain a simple formula (in the sense of a product of semigroups)

$$P_\tau^P = P_\tau^N P_\tau^M \tag{8}$$

where the upper index of the heat kernel denotes the manifold of its definition.

Hence

$$\mathcal{G}^m(X,X') = \frac{1}{2}\int_0^\infty d\tau \exp(-\frac{1}{2}m^2\tau)P_\tau^N(x,x')P_\tau^M(y,y') \tag{9}$$

The formula (9) is useful if we have a reliable approximation for the heat kernels on $\mathcal{N}$ and $\mathcal{M}$. We could conclude from eq.(9) (using the Schwinger-DeWitt asymptotic expansion) that if X is close to X' then $\mathcal{G} \simeq (s_N^2(x,x') + s_M^2(y,y'))^{-\frac{D}{2}+1}$, where s_M denotes the geodesic distance on $\mathcal{M}$. We are interested in the behaviour of the Green functions when $s_N >> s_M$. For such a purpose the formula (9) does not seem useful. We apply eigenfunction expansions of the heat kernels in the next section.

The special case of $D-d=2$ and $v=1$ can be studied in more detail. We choose the isometric coordinates with $g_{ab} = \delta_{ab}a^2$. In such a case eq.(4) reads

$$\Big(a^2(x)(-\triangle_M + m^2) - \triangle_2\Big)\mathcal{G}^m = h_M^{-\frac{1}{2}}\delta(X-X') \tag{10}$$

where $\triangle_2$ is the Laplacian on R^2.

3 Eigenfunction expansions

We assume in this section that $\mathcal{M}$ is a compact manifold without a boundary. Then, $-\triangle_M$ has a complete discrete set of orthonormal eigenfunctions [9]

$$-\triangle_M u_k = \epsilon_k u_k \tag{11}$$

satisfying the completeness relation

$$\sum_k \overline{u}_k(y)u_k(y') = h_M^{-\frac{1}{2}}\delta(y-y') \tag{12}$$

Let us note that 1 is an eigenfunction (11) with the eigenvalue 0 (we use the normalization $\int dy\sqrt{h_M} = 1$). Then, (distinguishing the zero mode) we can expand the heat kernel in eigenfunctions

$$P_\tau^M(y,y') = 1 + \sum_{k\neq 0}\exp(-\frac{1}{2}\epsilon_k\tau)\overline{u}_k(y)u_k(y') \tag{13}$$

For the $\mathcal{N}$ part of $\mathcal{P}$ we consider the eigenvalue problem in $L^2(dx)$ suggested by eq.(4)

$$\mathcal{A}_k\phi_E^k = \Big(v^{d-2}\sqrt{g}\omega_k^2 - \partial_a g^{ab}v^d\sqrt{g}\partial_b\Big)\phi_E^k = E_k\phi_E^k \tag{14}$$

where

$$\omega_k^2 = \epsilon_k + m^2 \tag{15}$$

The eigenfunctions satisfy the completeness relation

$$\sum_E \overline{\phi}_E^k(x)\phi_E^k(x') = \delta(x-x')$$

where the sum must be replaced by an integral if the spectrum of $\mathcal{A}_k$ in eq.(14) is continuous.

We expand the Green function in eigenfunctions u_k of the Laplace-Beltrami operator $\triangle_M$

$$\mathcal{G}^m(X,X') \equiv \textstyle\sum_k \mathcal{G}_k^m(X,X') = \sum_k g_k^m(x,x')\overline{u}_k(y)u_k(y') \tag{16}$$

Then, g_k is expanded in the eigenfunctions (14)

$$g_k^m(x,x') = \sum_E E_k^{-1}\overline{\phi}_E^k(x)\phi_E^k(x') \tag{17}$$

g_k^m is a solution of the equation

$$\mathcal{A}_k g_k^m(x,x') = \delta(x-x') \tag{18}$$

1 is an eigenfunction of $-\triangle_M$ with the eigenvalue 0. If we subtract the zero mode $\mathcal{G}_0^m$ (corresponding to $u_0 = 1$) from $\mathcal{G}^m$ then

$$\mathcal{G}^m(X,X') - \mathcal{G}_0^m(X,X') = \textstyle\sum_{E,k>0} E_k^{-1}\overline{\phi}_E^k(x)\phi_E^k(x')\overline{u}_k(y)u_k(y) \tag{19}$$

with

$$\mathcal{G}_0^m(X,X')=G^m(x,x') \tag{20}$$

where G^m is a solution of the equation

$$\Big(-\partial_a g^{ab}v^d g^{\frac{1}{2}}\partial_b+m^2v^{d-2}g^{\frac{1}{2}}\Big)G^m=\delta(x-x') \tag{21}$$

We investigate in this paper whether $\mathcal{G}^m$ can be approximated by $\mathcal{G}_0^m$,i.e., by the Green function G^m on $\mathcal{N}$. Such an approximation cannot be true for small distances because if the metric tensor is a regular function then the singularity of the Green function depends on the dimension of the space-time (see the discussion in sec.7). We expect that the approximation makes sense for large distances in $\mathcal{N}$. It can be seen that a decay of eigenfunctions ϕ_E^k is sufficient for a disappearance of each term on the rhs of eq.(19) at large x(this is not a necessary condition as we show soon). The eigenfunctions ϕ_E^k are localized if the spectrum of the operators $\mathcal{A}_k$ (14) is discrete. We can estimate the decay of eigenfunctions ϕ_E^k applying the eikonal (WKB) approximation to eq.(14). This means that we write $\phi^k=\exp(-\omega_k W)$ assuming that W is growing uniformly in each direction for large distances. Then, in the leading order we obtain for large x the equation

$$1=v^2g^{ab}\partial_aW\partial_bW \tag{22}$$

We obtain an exponential localization (increasing with the eigenvalue ϵ_k) of ϕ_E^k if $v^{-2}g_{ab}$ is uniformly growing to infinity for large distances. Let us note that $v^{-2}g_{ab}$ is the metric on $\mathcal{N}$ related to the one of eq.(1) by a conformal transformation. The growth of $\tilde{g}_{ab}=v^{-2}g_{ab}$ means that the volume element $\int dx\sqrt{\tilde{g}}$ is infinite. Such a property could be used to characterize the manifolds $\mathcal{P}$ whose Green function is dominated by the zero mode. Our rough arguments need a confirmation by a mathematical theory of the eigenvalue problems of second order differential operators (see [10] for some partial results). The sum over eigenvalues ϵ_k (the rhs of eq.(19)) will be discussed in sec.5.

In the special case (10) when $v=1$ and $D-d=2$ the eigenvalue equation (14) reduces to the well-known problem of quantum mechanics

$$\mathcal{A}\phi_E^k=(-\triangle_2+\omega_k^2a^2(x))\phi_E^k=E_k\phi_E^k \tag{23}$$

Here, $\triangle_2$ denotes the two-dimensional Laplacian. In this special case we have simple criteria for the discreteness of the spectrum of $\mathcal{A}$. If a^2 is growing uniformly in all directions then the spectrum of $\mathcal{A}$ is discrete and the eigenfunctions are localized (see [11] for a precise formulation and proofs).The eikonal approximation reads

$$\nabla W\nabla W=a^2(x) \tag{24}$$

Hence, $|\nabla W(x)|$ is growing like $a(x)$. The decay of eigenfunctions derived from eq.(24) is in agreement with exact results [11]. If the eigenfunctions ϕ_E^k decay for large x then the volume of $\mathcal{N}$, equal to $\int dxa^2(x)$, is infinite ($\mathcal{N}$ is not compact).

The localization of eigenfunctions $\phi^k_E(x)$ is not necessary for a decrease of $\mathcal{G}^m - \mathcal{G}^m_0$. Let us consider the simplest case of a non-compact $\mathcal{N} = R^{D-d}$ with $v = 1$ and $g_{ab} = \delta_{ab}$. Then, $\mathcal{A}_k$ has a continuous spectrum, its eigenfunctions are not localized, but

$$\begin{aligned}&(\mathcal{G}^m - \mathcal{G}^m_0)(X, X')\\ &= \textstyle\sum_{k\neq 0} \int_0^\infty d\tau \exp(-\frac{1}{2}\omega_k^2\tau - \frac{1}{2\tau}(x - x')^2)(2\pi\tau)^{-\frac{D-d}{2}}\overline{u}_k(y)u_k(y')\\ &\equiv \textstyle\sum_{k\neq 0} g^m_k(x - x')\overline{u}_k(y)u_k(y')\end{aligned} \tag{25}$$

where

$$\mathcal{G}^m_0(x, x') = g^m_0(x, x') = \textstyle\int_0^\infty d\tau \exp(-\frac{1}{2}m^2\tau - \frac{1}{2\tau}(x - x')^2)(2\pi\tau)^{-\frac{D-d}{2}} \tag{26}$$

and

$$g^m_k(x - x') = 2(2\pi)^{\nu-1}|x - x'|^\nu \omega_k^{-\nu} K_\nu(\omega_k|x - x'|)$$

with $\nu = -\frac{D-d}{2} + 1$, where K_ν is the modified Bessel function of the third kind [12]. From the asymptotic expansion of K_ν it follows that (for any $m^2 \geq 0$) each term on the rhs of eq.(25) is decaying exponentially for large $|x-x'|$. The sum on the rhs of eq.(26) will be estimated in sec.5.

If $\mathcal{N}$ is a compact manifold without a boundary then the spectrum λ_n of the Laplace-Beltrami operator $\triangle_N$ on $\mathcal{N}$ is discrete

$$-\triangle_N\psi_n = \lambda_n\psi_n \tag{27}$$

In such a case

$$P^N_\tau(x, x') = 1 + \sum_{n\neq 0} \exp(-\frac{1}{2}\lambda_n\tau)\overline{\psi}_n(x)\psi_n(x') \tag{28}$$

Hence, if $v = 1$ then from eq.(9)

$$(\mathcal{G}^m - \mathcal{G}^m_0)(X, X') = \textstyle\sum_{k>0} g_k(x, x')\overline{u}_k(y)u_k(y') \tag{29}$$

where

$$g^m_k(x, x') = \sum_n(\lambda_n + \epsilon_k + m^2)^{-1}\overline{\psi}_n(x)\psi_n(x') \tag{30}$$

and

$$\mathcal{G}^m_0(x, x') = G^m(x, x') = \textstyle\sum_n(\lambda_n + m^2)^{-1}\overline{\psi}_n(x)\psi_n(x') \tag{31}$$

is the Green function (3) on $\mathcal{N}$ (solving eq.(21) for $v = 1$).

However, if $\mathcal{N}$ is compact then there is no reason to neglect the rhs of eq.(29).

4 A two-dimensional manifold $\mathcal{N}$ with a Killing vector

We discuss in this section in more detail the product manifold $\mathcal{P} = \mathcal{N} \times \mathcal{M}$ ($v = 1$) when the two-dimensional manifold $\mathcal{N}$ has a symmetry generated by a Killing vector K. In an adapted system of coordinates such that $K = \partial_1$ the metric can be written in the form

$$ds^2 = dx_0^2 + a_1^2(x_0)dx_1^2 + \sum_{jk} h_{jk}(y)dy^j dy^k \tag{32}$$

In such a case the equation for the Green function reads

$$-(\partial_0 a_1 \partial_0 + a_1^{-1}\partial_1^2 + a_1 \triangle_M - m^2 a_1)\mathcal{G}^m = h_M^{-\frac{1}{2}}\delta \tag{33}$$

We can write the metric in an equivalent form. Let

$$\hat{x}_0 = \int dx_0 (a_1(x_0))^{-1} \tag{34}$$

Then,

$$ds^2 = a_1^2(\hat{x}_0)\Big(d\hat{x}_0^2 + dx_1^2\Big) + ds_M^2 \tag{35}$$

where ds_M^2 is the metric on $\mathcal{M}$. In the new coordinates

$$(-\hat{\partial}_0^2 - \partial_1^2 - a_1^2\triangle_M + m^2 a_1^2)\mathcal{G}^m = h_M^{-\frac{1}{2}}\hat{\delta} \tag{36}$$

where $\hat{\delta} = \delta(\hat{x} - \hat{x}')$ depends on $\hat{x}$ variables.

As an example, let $a_1(x_0) = x_0$ then

$$\hat{x}_0 = \ln(x_0) \tag{37}$$

Hence, eq.(36) takes the form

$$(-\hat{\partial}_0^2 - \partial_1^2 - \exp(2\hat{x}_0)(\triangle_M - m^2))\mathcal{G}^m = h_M^{-\frac{1}{2}}\hat{\delta} \tag{38}$$

In spite of a vanishing at zero (in the original x_0 coordinate) it can be checked by means of a calculation of the curvature tensor R that R is a continuous function (if $\mathcal{M} = R^n$ then the formula for the curvature in Bianchi type space-times derived in [13][14] gives $R = 0$). We have discussed the model (38) in detail in [2]. It has been shown that the model serves as an approximation to the Green function on a space-time with the bifurcate Killing horizon. If $\mathcal{M} = R^d$ then eq.(38) defines the Green function on the Euclidean version of the Rindler space.

An interesting class of models results from a choice of a metric which has a power-like singularity at x_0 when approaching $x_0 = 0$. We may choose $a_1(x_0)^2 =$

$\alpha|x_0|^{2\gamma}$ (the curvature tends to infinity at the singularity if $\gamma \neq 1$). Then, in the coordinates (34) we have

$$a_1(x_0)^2 = |\hat{x}_0|^{\frac{2\gamma}{1-\gamma}} \tag{39}$$

in eqs.(35)-(36).

We apply the eigenfunction expansion (23) of sec.3 to the case when a_1 depends only on x_0. We consider eq.(36) (we omit the hat over x_0). We expand the Green function in a complete set of orthonormal eigenfunctions of the one-dimensional quantum mechanical problem

$$(-\partial_0^2 + \omega_k^2 a_1(x_0)^2)\phi_n^k = \lambda_n(k)^2 \phi_n^k \tag{40}$$

where ω_k is defined in eq.(15) and

$$\sum_n \overline{\phi}_n^k(x_0)\phi_n^k(x_0') = \delta(x_0 - x_0') \tag{41}$$

Then, a solution of the equation for the Green function (10) has an expansion

$$\begin{aligned} &\mathcal{G}^m(X, X') \\ &= \pi^{-1} \textstyle\sum_{k,n} \overline{u}_k(y) u_k(y') \overline{\phi}_n^k(x_0)\phi_n^k(x_0')\lambda_n(k)^{-1} \exp(-\lambda_n(k)|x_1 - x_1'|) \end{aligned} \tag{42}$$

The formula (42) follows from eqs.(16)-(17) if we write

$$\phi_E^k(x_0, x_1) = \exp(ip_1 x_1)\phi_n^k(x_0)$$

with $E_k = p_1^2 + \lambda_n(k)^2$ and

$$\sum_E = \int dp_1 \sum_n$$

Then, the integral over p_1 in eq.(17) leads to eq.(42).

The eigenfunctions ϕ^k are decaying exponentially if a_1 is growing at infinity, as can be seen, e.g., from the WKB approximation (see [11] for rigorous results)

$$\phi_n^k(x) \simeq \exp(-\omega_k \int dx a_1(x)) \tag{43}$$

Hence, each term in the expansion (42) is decaying exponentially. Note that according to eqs.(22),(24) and (43) the terms with larger eigenvalues ϵ_k are decaying faster then the ones with the lower eigenvalue.

5 The correction to the contribution of the zero mode

We expect that in general (for a non-compact manifold $\mathcal{N}$) the difference $\mathcal{G}^m - \mathcal{G}_0^m$ is negligible for large distances on $\mathcal{N}$. First, we must estimate the Green functions g_k^m (18) of the second order differential operators $\mathcal{A}_k$(depending on ϵ_k) on $\mathcal{N}$ for large distances. We write

$$g_k^m(x,x') = \exp(-\omega_k W(x,x')) \tag{44}$$

Assuming that W is growing uniformly in each direction we obtain in the leading order for large distances eq.(22) for W. We recognize eq.(22) as an equation for a geodesic distance s_N^v on the manifold $\mathcal{N}$ with the metric $v^{-2}g_{ab}$ [15]. Hence, the geodesic distance $W(x,x') = s_N^v(x,x')$ is the solution of eq.(22) which is symmetric under the exchange of the points and satisfies the boundary condition $W(x,x) = 0$. We insert the approximate solutions g_k^m (44) (for some rigorous results on the large distance behaviour of Green functions of second order differential operators see [10][16]) into the sum (19) over eigenvalues and eigenfunctions of $-\triangle_M$. We approximate the sum over large eigenvalues by an integral (Weyl approximation) assuming $|u_k(y)| \leq C$. Then

$$\begin{aligned} &|(\mathcal{G}^m - \mathcal{G}_0^m(X,X')| \\ &\leq A_1 \sum_{k<\Lambda} \exp(-\omega_k s_N^v(x,x'))|u_k(y)||u_k(y')| + A_2 \int_{|\mathbf{k}|>\Lambda} d\mathbf{k} \exp(-\omega_k s_N^v(x,x')) \end{aligned} \tag{45}$$

where in the integral over $\mathbf{k}$ we set $\omega_k = \sqrt{\mathbf{k}^2 + m^2}$. The integral over $\mathbf{k}$ is decreasing exponentially as a function of s_N.

We make the estimate precise in the simple model of $\mathcal{N} = R^{D-d}$ (eq.(25)). We apply Weyl theory [17] saying that for large eigenvalues the sum over eigenvalues of the Laplace-Beltrami operator on a d-dimensional compact manifold $\mathcal{M}$ can be approximated by a d-dimensional integral with $\epsilon_k \simeq \mathbf{k}^2$. If additionally we assume $|u_k| \leq C$ then in eq.(25)

$$|(\mathcal{G}^m - \mathcal{G}_0^m)(X,X')| \leq A_1 \sum_{k<\Lambda} |c_k(x-x')||u_k(y)||u_k(y')| + R_\Lambda(x-x') \tag{46}$$

where from the Weyl approximation

$$\begin{aligned} R_\Lambda(x-x') &= C^2 \int_0^\infty d\tau \int_{|\mathbf{k}|>\Lambda} d\mathbf{k} \exp(-\tfrac{1}{2}(\mathbf{k}^2+m^2)\tau - \tfrac{1}{2\tau}(x-x')^2)(2\pi\tau)^{-\frac{D-d}{2}} \\ &= A \int_0^\infty d\tau \tau^{-\frac{d}{2}} \Gamma(\tfrac{d}{2}, \tfrac{1}{2}\tau\Lambda^2) \exp(-\tfrac{1}{2}m^2\tau - \tfrac{1}{2\tau}(x-x')^2)(2\pi\tau)^{-\frac{D-d}{2}} \end{aligned} \tag{47}$$

where Γ denotes the incomplete gamma function [12]. From the asymptotic expansion of Γ we obtain that

$$R_\Lambda(x-x') \simeq \exp(-\sqrt{\Lambda^2+m^2}|x-x'|) \tag{48}$$

for large $|x-x'|$. Hence, we can conclude that $\mathcal{G}^m - \mathcal{G}_0^m$ for large $|x-x'|$ is decreasing as $\exp(-\sqrt{m^2+\epsilon}|x-x'|)$, where ϵ is the lowest non-zero eigenvalue of $-\triangle_M$. The decay of $|(\mathcal{G}^m - \mathcal{G}_0^m)(X,X')|$ for large distances is determined by the first term on the rhs of eq.(46).

We repeat the estimates in the model (25) without any reference to the Weyl approximation in the simplest case of $\mathcal{M} = S^\beta$ where S^β is the circle of radius

β (then the spectrum of $-\triangle_M$ is known). The method of an explicit sum over eigenvalues is simple if $m = 0$. Then, from eqs.(9) and (17)

$$\begin{aligned} &\mathcal{G}^0(X,X') \\ &= (2\pi\beta)^{-1} \int_0^\infty d\tau (2\pi\tau)^{-\frac{D-1}{2}} \exp(-\tfrac{1}{2\tau}(x-x')^2) \sum_k \exp(-\tfrac{\tau}{2}(\tfrac{k}{\beta})^2) \exp(i\tfrac{k}{\beta}(y-y')) \\ &= \int_0^\infty d\tau (2\pi\tau)^{-\frac{D}{2}} \exp(-\tfrac{1}{2\tau}(x-x')^2) \sum_k \exp(-\tfrac{1}{2\tau}(y-y'-2\pi\beta k)^2) \end{aligned} \tag{49}$$

Performing the integral over τ we obtain

$$\mathcal{G}^0(X,X') = (2\pi)^{-D+2}\Gamma(\tfrac{D}{2}-1) \sum_k ((x-x')^2 + (y-y'-2\pi\beta k)^2)^{-\frac{D}{2}+1} \tag{50}$$

In order to perform the sum we apply the formula

$$\begin{aligned} &\sum_k (\sigma^2 + (y-y'-2\pi\beta k)^2)^{-1} \\ &= \tfrac{1}{4\beta\sigma}\Big(\coth(\tfrac{1}{2\beta}(\sigma + i(y-y')) + \coth(\tfrac{1}{2\beta}(\sigma - i(y-y'))\Big) \end{aligned} \tag{51}$$

The formula (51) can be applied directly to eq.(50) with $\sigma^2 = (x-x')^2$ if $D = 4$. When D is even and bigger then 4 then we differentiate eq.(51) over σ^2 and subsequently apply to the sum (50). If $D = 2n+1$ is odd then we have to use differentiation $n-1$ times together with an integration in order to perform the sum in eq.(50).

We show this procedure for $D = 3$. This dimension is relevant for the model (25) of sec.3 and in models of sec.4. We use the integral

$$\int_0^\infty dr (r^2 + a^2)^{-1} = \pi a^{-1} \tag{52}$$

in order to represent the Green function in $D = 3$ dimensions in the form

$$\begin{aligned} &\mathcal{G}^0(X,X') \\ &= 2^{-\frac{3}{2}}\pi^{-2}\beta^{-1} \int_0^\infty dr \tfrac{1}{4\sigma}\Big(\coth(\tfrac{1}{2\beta}(\sigma + i(y-y')) + \coth(\tfrac{1}{2\beta}(\sigma - i(y-y'))\Big) \end{aligned} \tag{53}$$

where $\sigma^2 = r^2 + (x_0 - x_0')^2 + (x_1 - x_1')^2$. Then,

$$\begin{aligned} &(\mathcal{G}^0 - \mathcal{G}^0_0)(X,X') \\ &= 2^{-\frac{3}{2}}\pi^{-2}\beta^{-1} \int_0^\infty dr \tfrac{1}{4\sigma}\Big(\coth(\tfrac{1}{2\beta}(\sigma + i(y-y'))) - 1 + \coth(\tfrac{1}{2\beta}(\sigma - i(y-y')) - 1\Big) \end{aligned} \tag{54}$$

where $\mathcal{G}^0_0 = G^0$ is defined in eq.(26) with $D - d = 2$ (for $m = 0$ and $D - d = 2$ the integral (26) is divergent at large τ; this is the well-known infrared problem for massless scalar fields, it can be avoided by a choice of test functions for the smeared out fields with no support at the zero momentum, in such a case the integral (26) is defined as a logarithm of the distance).

Applying the formula

$$\coth v - 1 = 2\exp(-2v)(1-\exp(-2v))^{-1}$$

we bound the integral on the rhs of eq.(54) by

$$|(\mathcal{G}^0-\mathcal{G}^0_0)(X,X')| \leq A\int_0^a dr\sigma^{-1}\exp(-2\sigma) = AK_0\Big(\frac{1}{\beta}\sqrt{(x_0-x_0')^2+(x_1-x_1')^2}\Big)$$

for large $(x_0-x_0')^2+(x_1-x_1')^2$, where K_ν denotes the modified Bessel function of the third kind [12], which is exponentially decreasing for large arguments.

We conclude from eq.(54) that the logarithmic two-dimensional propagator well approximates the three dimensional propagator for distances $(x_0-x_0')^2+(x_1-x_1')^2 >> \beta^2$. Applying eqs.(50)-(51) we could confirm for any dimension $D-d$ the result following from eqs.(46)-(48) that the difference $\mathcal{G}^0-\mathcal{G}^0_0$ is exponentially small for large distances (the length scale is β which is equal to the radius of the circle S^β or in other words it is the square root of the inverse of the lowest non-zero eigenvalue of $-\triangle_M$). For $D>3$ the Green function $\mathcal{G}$ would be approximated by $\mathcal{G}^0_0 \simeq |x-x'|^{-D+3}$ (by $\log|x-x'|$ in $D=3$). The high temperature limit of interacting field theories is discussed in [18][19].

6 Quantum free fields on the product manifold

We introduce now a free Euclidean field as a random field with the two-point correlation function equal to the Green function (see [7]). The Green function (17) defines the Gaussian Euclidean field

$$\Phi(x,y) \equiv \sum_k \Phi_k = \sum_k \chi_k(x)u_k(y) \tag{55}$$

where

$$\langle\chi_k(x)\chi_r(x')\rangle = \delta_{kr}g^m_k(x,x') \tag{56}$$

g^m_k as a Green function of the second order differential operator is non-negative. This is a positive definite bilinear form. Hence, it defines Gaussian Euclidean field ϕ_k on $\mathcal{N}$. In the example (25) of $\mathcal{N}=R^{D-d}$ we have $g^m_k=(-\triangle+\omega_k^2)^{-1}$.Then, ϕ^m_k is the Euclidean free field on R^{D-d} with the mass ω_k (ϕ^m_0 has the mass m). In an analytic continuation of the model (25) to the Minkowski space χ_k becomes the free quantum field with the mass ω_k on the $D-d$ dimensional Minkowski space.

The models of sec.4 have an expansion

$$\begin{aligned}&\Phi(x_0,x_1,y) \equiv \textstyle\sum_k \Phi_k = \int dp_1 \exp(ip_1x_1)\sum_{k,n} a_k(p_1,n)\phi^k_n(x_0)u_k(y)\\ &= \Phi_0(x_0,x_1)+\textstyle\sum_{k>0}\Phi_k(x_0,x_1,y)\end{aligned} \tag{57}$$

where

$$\langle\overline{a}_k(p_1,n)a_{k'}(p_1',n')\rangle = \delta(p_1-p_1')\delta_{nn'}\delta_{kk'}(p_1^2+\lambda_n(k)^2)^{-1} \tag{58}$$

We have

$$\langle\Phi_0(x_0,x_1)\Phi_0(x_0',x_1')\rangle = \mathcal{G}^m_0(x,x') = \int_0^\infty d\tau\exp(-\tfrac{1}{2}m^2\tau-\tfrac{1}{2\tau}(x-x')^2)(2\pi\tau)^{-1} \tag{59}$$

and for $k > 0$

$$\begin{aligned}&\langle\overline{\Phi}_k(x_0,x_1,y)\Phi_k(x_0',x_1',y')\rangle\\ &=\frac{1}{\pi}\sum_n \lambda_n(k)^{-1}\exp(-\lambda_n(k)|x_1-x_1'|)\overline{\phi}_n^k(x_0)\phi_n^k(x_0')\overline{u}_k(y)u_k(y')\end{aligned}\tag{60}$$

We can continue analytically the Green functions (60) of the model (57) into imaginary values of x_1 (then x_1 plays the role of time; the analytic continuation follows from the positivity of the Green function (60) under a reflection of x_1 [7])

$$\begin{aligned}&\mathcal{G}^m(x_0,ix_1,y,ix_1',y')\\ &=\frac{1}{\pi}\sum_{k,n} \lambda_n(k)^{-1}\exp(i\lambda_n(k)(x_1-x_1'))\overline{\phi}_n^k(x_0)\phi_n^k(x_0')\overline{u}_k(y)u_k(y')\end{aligned}\tag{61}$$

Such Green functions result from the quantization in the Fock space

$$\begin{aligned}&\Phi(x_0,x_1,y)=\sum_{k,n}\exp(-i\lambda_n(k)x_1)a_k(n)\phi_n^k(x_0)u_k(y)\\ &+\sum_{k,n}\exp(i\lambda_n(k)x_1)a_k^+(n)\phi_n^k(x_0)\overline{u}_k(y)\end{aligned}\tag{62}$$

where

$$[a_k(n),a_{k'}^+(n')]=\frac{1}{\pi}\delta_{nn'}\delta_{kk'}\lambda_n(k)^{-1}\tag{63}$$

are the creation and annihilation operators.

In order to define a quantum field from the Euclidean field (60) with another time we would need the reflection positivity in another direction . Various analytic continuations are possible on the same manifold $\mathcal{N}$. As an example, if $\mathcal{N}$ is the hyperbolic space then the Euclidean field on $\mathcal{N}$ can be analytically continued either to a quantum field on DeSitter space or to the one defined on the AntiDeSitter space [7]. The model (38) of sec.4 ($a_1 \simeq x_0^2$) with $\mathcal{M} = R^d$ as the Euclidean version of the Rindler space has an analytic continuation to the Rindler space or to the Milne space. The model is also related by a conformal transformation to the direct product of the real line and the hyperbolic space (the models and their analytic continuations are discussed in [20] and in the references cited there).

In general, if we are able to define a quantum field with the Green function defined by the analytic continuation of g_k^m (18) (for every k) then we can define the quantum field on an analytic continuation of $\mathcal{P}$ by means of the expansion (55).

7 Discussion

We have investigated some aspects of quantum free fields on product manifolds $\mathcal{P} = \mathcal{N} \times \mathcal{M}$. Such studies have a long history. It is an old argument that on a compact manifold $\mathcal{M}$ the heavy modes (KK-modes) decouple from the zero mode (defining the dimensional reduction). In this paper we have studied the problem in the configuration space. An approximation of quantum field theory by a lower dimensional one is not trivial because the theories in various dimensions have different short distance behaviour. The dimensional reduction of the Kaluza-Klein type can make sense only above a certain length scale determined by the size of the compact manifold (or equivalently by the inverse of the square root of the lowest non-zero eigenvalue of $-\triangle_M$). In sec.5 we have estimated the difference $\mathcal{G}-\mathcal{G}_0$, where $\mathcal{G}_0$ is the $D-d$ dimensional Green function. We have shown in some models that there is no dimensional reduction for a non-compact $\mathcal{M}$ [2]. Nevertheless, a study of Green functions with non-compact manifolds $\mathcal{M}$ can lead to some unexpected results. The conventional behaviour $|x-x'|^{-D+2}$ in D dimensions of the Green functions of quantum massless fields can fail on a manifold. The Green functions of the model of sec.4 can be more singular than it would follow from the dimension-dependent index $D-2$. A change of the short distance behaviour has been shown in the brane model of Dvali et al [5]. Their model can be formulated in the framework of sec.2 with $D-d=1$. The equation for the Green function reads

$$(-\partial_0^2+a^2(x_0)(-\triangle+m^2))\mathcal{G}^m=\delta \tag{64}$$

The Fourier transform of $\mathcal{G}$ in $D-1$ variables satisfies an equation for the quantum mechanical Green function with the potential $V=\mathbf{p}^2a^2(x_0)$. We have studied solutions of the equations for Green functions in [21] by means of path integrals. The short distance behaviour depends on the singularity of $a^2(x_0)$ for $x_0\rightarrow 0$. In the model of ref.[5]

$$a^2(x_0)=\alpha+\beta\delta(x_0) \tag{65}$$

It follows from [5][21] that $\mathcal{G}(0,y,0,y')\simeq|y-y'|^{-D+3}$. This is the behaviour of the Green function in $D-1$ dimensions. Such a behaviour of the Green function at short distances may be called a dimensional reduction in spite of the continuous spectrum of $\triangle$ on $\mathcal{M}$.

Let us still consider the models with a singular metric of sec.4 in view of the results of [21]. The equation for the Green function reads

$$(-\hat{\partial}_0^2-\partial_1^2+|\hat{x}_0|^{\frac{2\gamma}{1-\gamma}}(-\triangle+m^2))\mathcal{G}^m=\hat{\delta} \tag{66}$$

where $\hat{x}_0$ has been called η in [21]. Applying the methods and results of ref.[21] we obtain

$$\mathcal{G}(0,x_1,0;0,x_1',0)\simeq|x_1-x_1'|^{\frac{-D+2}{1-\gamma}} \tag{67}$$

and

$$\mathcal{G}(0,0,y;0,0,y')\simeq|y-y'|^{-D+2} \tag{68}$$

for small distances. Hence, the Green functions in the x_1 direction are more singular (if $\gamma>0$)than the ones in R^D, whereas the singularity in the y direction is the same as the one in the D dimensional free field theory on a flat space.

In the model of Dvali et al [5] the long distance behaviour of massless fields is the same as the one in D dimensions. The dimensional reduction discussed in the present paper concerns the long distance behaviour. The short distance behaviour can be different from the canonical one (with the index $D-2$) only if the metric is singular.

An analytic continuation of Green functions of the models of sec.4 with a Killing vector on $\mathcal{N}$ as discussed in sec.6 follows from the general theory of a quantization of the symmetry generated by the Killing vector formulated in [7]. In the system of coordinates adapted to this Killing vector the analytic continuation concerns the coordinate x_1. It gives the Hilbert space, the Hamiltonian and a unitary evolution. There can be more reflection symmetries on a manifold. The Riemannian manifolds of sec.4 are invariant under the reflection $x_0 \to -x_0$. However, a reflection positivity (with respect to x_0), necessary for a definition of the quantum field (with x_0 as an imaginary time), remains unclear. The reflection positivity may have a physical meaning. Tunnelling between different topologies of space-time may require a reflection invariant Riemannian manifold [22]. Models with a prior to the Big Bang evolution [23][24] seem to require a Euclidean version which is invariant under a time reflection.

References

[1] Modern Kaluza-Klein Theories, Th. Appelquist, A. Chodos and P.G.O. Freund,eds., Addison-Wesley Publishing,1987,

C.A. Orzalesi, Fortschr.Phys.**29**,413(1981)

[2] Z. Haba, arXiv:0708.1742

[3] T. Padmanabhan, Mod.Phys.Lett.**A17**,923(2002), arXiv:gr-qc/0202078

T. Padmanabhan, Phys.Rep.**406**,49(2005)

[4] L.Randall, R.Sundrum, Phys.Rev.Lett. **83**,3370(1999);

Phys.Rev.Lett.**83**,4690(1999)

[5] G.Dvali,G.Gabadadze and M.Poratti, Phys.Lett.**485B**,208(2000)

G. Dvali and G. Gabadadze, Phys.Rev.**D63**,065007(2001)

[6] J. Dimock, Rev.Math.Phys.**16**,243(2004)

[7] A.Jaffe and G. Ritter, Commun.Math.Phys,**270**,545(2007)

[8] J. Choi, J.Math.Phys.**41**,8163(2000);

J. Choi and S.-T.Hong,J.Math.Phys.**45**,642(2004)

[9] I. Chavel, Eigenvalues in Riemannian Geometry, Academic Press, New York,1984

[10] S. Agmon, Lectures on Exponential Decay of Solutions of Second-Order Elliptic Equations, Princeton Univ.Press, Princeton, New Jersey,1982

[11] M. Reed and B. Simon, Methods of Mathematical Physics, Academic Press, New York,1979

[12] I.S. Gradstein and I.M.Ryzhik, Table of Integrals,Series and Products, Academic Press, New York,1962

[13] S.A. Fulling, L. Parker and B.L. Hu, Phys.Rev.**D10**,3905(1974)

[14] N. D. Birrell and P.C.W. Davis, Quantum Fields in Curved Space, Cambridge University Press,1982

[15] DeWitt B.S. and Brehme R.W., Ann.Phys.**9**,220(1960)

[16] E.B. Davis and E.M. Harrell , Journ.Diff.Eqs.**66**,165(1987)

[17] M.E. Taylor,Pseudodifferential 0perators, Princeton University Press,1981

[18] L. Dolan and R. Jackiw, Phys.Rev.**D9**,3320(1974)

[19] Th. Appelquist and R.D. Pisarski, Phys.Rev. **D23**,2305(1981)

[20] Z. Haba, Journ.Phys.**A40**,8535(2007)

[21] Z. Haba, Journ.Math.Phys.**46**,042301(2005)
Mod.Phys.Lett.**A32**,2411(2004)

[22] G.W. Gibbons and J.B. Hartle, Phys.Rev.**D42**,2458(1990)
G.W. Gibbons and H.-J.Pohle, Nucl.Phys.**B410**,117(1993)

[23] G. Veneziano, arXiv:hep-th/0002094

[24] J. Khoury, B.A. Ovrut, N. Seiberg, P.J. Steinhardt and N. Turok, arxiv:hep-th/0108187